



Data Collection Worksheet

Please Note: The Data Collection Worksheet (DCW) is a tool to aid integration of a PhenX protocol into a study. The PhenX DCW is not designed to be a data collection instrument. Investigators will need to decide the best way to collect data for the PhenX protocol in their study. Variables captured in the DCW, along with variable names and unique PhenX variable identifiers, are included in the PhenX Data Dictionary (DD) files.

The Tobacco, Alcohol, Prescription medication, and other Substance (TAPS) Tool Part 1

General Instructions:

The TAPS Tool Part 1 is a 4-item screening for tobacco use, alcohol use, prescription medication misuse, and illicit substance use in the past year. Question 2 should be answered only by males and Question 3 only by females. Each of the four multiple-choice items has five possible responses to choose from. Check the box to select your answer.

Segment:

Visit number:

1. In the PAST 12 MONTHS, how often have you used any tobacco product (for example, cigarettes, e-cigarettes, cigars, pipes, or smokeless tobacco)?

☐ Daily or Almost Daily

☐ Weekly

☐ Monthly

☐ Less Than Monthly

☐ Never

2. In the PAST 12 MONTHS, how often have you had 5 or more drinks containing alcohol in one day? One standard drink is about 1 small glass of wine (5 oz), 1 beer (12 oz), or 1 single shot of liquor. (Note: This question should only be answered by males).

☐ Daily or Almost Daily

☐ Weekly

☐ Monthly

☐ Less Than Monthly

☐ Never

3. In the PAST 12 MONTHS, how often have you had 4 or more drinks containing alcohol in one day? One standard drink is about 1 small glass of wine (5 oz), 1 beer (12 oz), or 1 single shot of liquor. (Note: This question should only be answered by females).

☐ Daily or Almost Daily

☐ Weekly

☐ Monthly

☐ Less Than Monthly

☐ Never

4. In the PAST 12 MONTHS, how often have you used any drugs including marijuana, cocaine or crack, heroin, methamphetamine (crystal meth), hallucinogens, ecstasy/MDMA?

☐ Daily or Almost Daily

☐ Weekly

☐ Monthly

☐ Less Than Monthly

☐ Never

5. In the PAST 12 MONTHS, how often have you used any prescription medications just for the feeling, more than prescribed or that were not prescribed for you? Prescription medications that may be used this way include: Opiate pain relievers (for example, OxyContin, Vicodin, Percocet, Methadone), Medications for anxiety or sleeping (for example, Xanax, Ativan, Klonopin), Medications for ADHD (for example, Adderall or Ritalin)

☐ Daily or Almost Daily

☐ Weekly

☐ Monthly

☐ Less Than Monthly

☐ Never

Protocol source: <https://www.phenxtoolkit.org/protocols/view/511601>